

Clinical Policy: Ponesimod (Ponvory)

Reference Number: CP.PHAR.537 Effective Date: 06.01.21 Last Review Date: 05.24 Line of Business: Commercial, HIM, Medicaid

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Ponesimod (Ponvory[®]) is a sphingosine 1-phosphate receptor modulator.

FDA Approved Indication(s)

Ponvory is indicated for the treatment of relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Ponvory is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Multiple Sclerosis (must meet all):
 - 1. Diagnosis of one of the following (a, b, or c):
 - a. Clinically isolated syndrome, and member is contraindicated to both, or has experienced clinically significant adverse effects to one, of the following at up to maximally indicated doses: an interferon-beta agent (Avonex[®], Betaseron[®]/Extavia^{®†}, Rebif[®], or Plegridy[®]), glatiramer (Copaxone[®], Glatopa[®]);
 - b. Relapsing-remitting MS, and failure of all of the following at up to maximally indicated doses, unless clinically significant adverse effects are experienced or all are contraindicated (i, ii, iii, and iv):
 - i. **Dimethyl fumarate** (generic Tecfidera[®]);
 - ii. **Teriflunomide** (generic Aubagio[®]);
 - iii. Fingolimod (Gilenya[®]);
 - iv. An **interferon-beta agent** (Avonex, Betaseron/Extavia[†], Rebif, or Plegridy) or **glatiramer** (Copaxone, Glatopa);

*Prior authorization may be required for all disease modifying therapies for MS †Betaseron is preferred for the Commercial and HIM lines of business; Extavia is preferred for the Medicaid line of business

- c. Secondary progressive MS;
- 2. Prescribed by or in consultation with a neurologist;
- 3. Age \geq 18 years;
- 4. Ponvory is not prescribed concurrently with other disease modifying therapies for MS (*see Appendix D*);

CLINICAL POLICY Ponesimod



- 5. Documentation of both baseline number of relapses per year and expanded disability status scale (EDSS) score;
- 6. Request meets the following:
 - a. Treatment initiation: Dose follows the 14-day titration schedule as outlined in Section V and does not exceed 1 tablet per day;
 - b. Treatment maintenance: Dose does not exceed (i and ii):
 - i. 20 mg per day;
 - ii. 1 tablet per day.

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

- A. Multiple Sclerosis (must meet all):
 - 1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
 - 2. Member meets one of the following (a or b):
 - a. If member has received < 1 year of total treatment: Member is responding positively to therapy;
 - b. If member has received ≥ 1 year of total treatment: Member meets one of the following (i, ii, iii, or iv):
 - i. Member has not had an increase in the number of relapses per year compared to baseline;
 - ii. Member has not had ≥ 2 new MRI-detected lesions;
 - iii. Member has not had an increase in EDSS score from baseline;

CLINICAL POLICY Ponesimod



- iv. Medical justification supports that member is responding positively to therapy;
- 3. Ponvory is not prescribed concurrently with other disease modifying therapies for MS (*see Appendix D*);
- 4. If request is for a dose increase, new dose does not exceed (a and b):
 - a. 20 mg per day;
 - b. 1 tablet per day.

Approval duration:

If member has received < 1 year of total treatment – up to a total of 12 months of treatment

If member has received ≥ 1 year of total treatment – 12 months

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents;
- **B.** Primary progressive MS.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key EDSS: expanded disability status scale FDA: Food and Drug Administration MS: multiple sclerosis



Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
teriflunomide (Aubagio [®])	7 mg or 14 mg PO QD	14 mg/day
Avonex [®] , Rebif [®] (interferon	Avonex: 30 mcg IM Q week	Avonex: 30 mcg/week
beta-1a)	<i>Rebif</i> : 22 mcg or 44 mcg SC TIW	Rebif: 44 mcg TIW
Betaseron [®] , Extavia [®]	250 mcg SC QOD	250 mg QOD
(interferon beta-1b)		
Plegridy [®] (peginterferon beta-	125 mcg SC Q2 weeks	125 mcg/2 weeks
1a)		_
glatiramer acetate	20 mg SC QD or 40 mg SC TIW	20 mg/day or 40 mg
(Copaxone [®] , Glatopa [®])		TIW
fingolimod (Gilenya [®])	0.5 mg PO QD	0.5 mg/day
dimethyl fumarate	120 mg PO BID for 7 days,	480 mg/day
(Tecfidera [®])	followed by 240 mg PO BID	

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): in the last 6 months, experienced myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III or IV heart failure; presence of Mobitz type II second-degree, third-degree atrioventricular block, sick sinus syndrome, or sino-atrial block, unless patient has a functioning pacemaker
- Boxed warning(s): none reported

Appendix D: General Information

- Disease-modifying therapies for MS are: glatiramer acetate (Copaxone[®], Glatopa[®]), interferon beta-1a (Avonex[®], Rebif[®]), interferon beta-1b (Betaseron[®], Extavia[®]), peginterferon beta-1a (Plegridy[®]), dimethyl fumarate (Tecfidera[®]), diroximel fumarate (Vumerity[®]), monomethyl fumarate (Bafiertam[™]), fingolimod (Gilenya[®], Tascenso ODT[™]), teriflunomide (Aubagio[®]), alemtuzumab (Lemtrada[®]), mitoxantrone (Novantrone[®]), natalizumab (Tysabri[®], and biosimilar Tyruko[®]), ocrelizumab (Ocrevus[®]), siponimod (Mayzent[®]), cladribine (Mavenclad[®]), ozanimod (Zeposia[®]), ofatumumab (Kesimpta[®]), ublituximab-xiiy (Briumvi[™]), and ponesimod (Ponvory[™]).
- Of the disease-modifying therapies for MS that are FDA-labeled for clinically isolated syndrome, only the interferon products, glatiramer, and teriflunomide have demonstrated any efficacy in decreasing the risk of conversion to MS compared to placebo. This is supported by the American Academy of Neurology 2018 MS guidelines.



V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose	
MS	Treatment initiation:	20 mg/day	
	• Days 1 and 2: 2 mg PO QD		
	• Days 3 and 4: 3 mg PO QD		
	• Days 5 and 6: 4 mg PO QD		
	• Day 7: 5 mg PO QD		
	• Day 8: 6 mg PO QD		
	• Day 9: 7 mg PO QD		
	• Day 10: 8 mg PO QD		
	• Day 11: 9 mg PO QD		
	• Day 12, 13, and 14: 10 mg PO QD		
	Treatment maintenance:		
	• Day 15 and thereafter: 20 mg PO QD		
	Missed doses:		
	• If fewer than 4 consecutive doses are missed:		
	• During titration: Resume treatment with the first		
	missed titration dose and resume the titration		
	schedule at that dose and titration day.		
	• During maintenance: Resume treatment with the maintenance dosage.		
	• If 4 or more consecutive doses are missed during		
	titration or maintenance:		
	• Treatment should be reinitiated with Day 1 of the		
	titration regimen (new starter pack).		

VI. Product Availability

- Tablets, starter pack: 2 mg, 3 mg, 4 mg, 5 mg, 6 mg, 7 mg, 8 mg, 9 mg, 10 mg
- Tablets, maintenance dose bottle: 20 mg

VII. References

- 1. Ponvory Prescribing Information. Titusville, NJ: Janssen Pharmaceuticals, Inc.; August 2023. Available at: https://www.ponvory.com. Accessed January 12, 2024.
- Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: disease-modifying therapies for adults with multiple sclerosis: report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. Neurology. 2018; 90(17): 777-788. Full guideline available at: https://www.aan.com/Guidelines/home/GetGuidelineContent/904. Reaffirmed on September 18, 2021.



Reviews, Revisions, and Approvals	Date	P&T Approval
		Date
Policy created	03.24.21	05.21
2Q 2022 annual review: no significant changes; clarified	02.08.22	05.22
interferon-beta product redirections for each line of business per		
SDC; references reviewed and updated.		
Template changes applied to other diagnoses/indications and	10.05.22	
continued therapy section.		
2Q 2023 annual review: no significant changes; to be inclusive of	01.31.23	05.23
members continuing therapy from a different benefit, revised		
continued approval duration to reference the duration of total		
treatment received rather than the number of re-authorizations;		
references reviewed and updated.		
Per August SDC, added generic references to Aubagio and Gilenya	08.22.23	11.23
redirections.		
2Q 2024 annual review: no significant changes; references	01.30.24	05.24
reviewed and updated.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan

CLINICAL POLICY Ponesimod



retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2021 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation.